

US009220590B2

(12) United States Patent

er (45) Date of Pate

(54) ACCOMMODATIVE INTRAOCULAR LENS AND METHOD OF IMPROVING ACCOMMODATION

(75) Inventor: Paul Marius Beer, Slingerlands, NY

(US)

(73) Assignee: Z Lens, LLC, Slingerlands, NY (US)

(*) Notice: Subject to any disclaimer, the term of this

patent is extended or adjusted under 35

U.S.C. 154(b) by 1163 days.

(21) Appl. No.: 13/157,703

(22) Filed: Jun. 10, 2011

(65) Prior Publication Data

US 2011/0307058 A1 Dec. 15, 2011

Related U.S. Application Data

- (60) Provisional application No. 61/353,273, filed on Jun. 10, 2010, provisional application No. 61/368,862, filed on Jul. 29, 2010.
- (51) **Int. Cl. A61F 2/16** (2006.01)

(56) References Cited

U.S. PATENT DOCUMENTS

3,975,779 A 8/1976 Richards et al. 4,373,218 A 2/1983 Schachar (10) Patent No.: US 9,220,590 B2 (45) Date of Patent: Dec. 29, 2015

4,463,457 A 8/1984 Kelman 4,527,294 A 7/1985 Heslin 4,534,069 A 8/1985 Kelman (Continued)

FOREIGN PATENT DOCUMENTS

GB 1583193 A 1/1981

OTHER PUBLICATIONS

International Search Report and Written Opinion mailed Dec. 4, 2013 in International Application No. PCT/US2013/039708 (18 pages).

(Continued)

Primary Examiner — Elizabeth Houston


Assistant Examiner — Joshua Levine

(74) Attorney, Agent, or Firm — Goodwin Procter LLP

(57) ABSTRACT

The present invention provides an accommodative intraocular lens (AIOL) system and method for improving accommodation with an intraocular lens. The method involves insertion into the capsular bag of a flexible optic holder comprising a plurality of haptics configured to allow the capsular bag to be sectioned at regular intervals following fusion of the capsular bag. The haptics of the optic holder are designed to allow maximum fusion of the anterior and posterior leaves of the capsular bag following placement of the optic holder in the capsular bag. Following introduction of the optic holder into the capsular bag, the natural or assisted process of fibrosis/ fusion of the capsular bag occurs, thereby sealing and securely capturing the haptics within the capsular bag. Subsequently, several cuts are made in the fibrotic capsular bag at intervals between haptics, allowing the haptics to move independently, thereby effectively restoring some of the flexibility that the capsule possessed prior to fibrosis and restoring some of the zonular force on the capsule.

20 Claims, 13 Drawing Sheets



US 9,220,590 B2 Page 2

(56)		Re	eferen	ces Cited	6,228,115			Hoffmann et al.
	ī	IS PAT	FENT	DOCUMENTS	6,241,777 6,251,312		6/2001 6/2001	Phan et al.
		J.D. 111.	LLIVI	DOCOMENTO	6,261,321		7/2001	Kellan
4	4,575,373 A	A 3	/1986	Johnson	6,280,449		8/2001	
	4,581,032 A			Grandon	6,299,641 6,302,912		10/2001 10/2001	
	4,581,033 <i>A</i>			Callahan	6,306,167			Bernau et al.
	4,666,446 <i>1</i> 4,685,922 <i>1</i>			Koziol et al. Peyman	6,358,280			Herrick
	4,693,717 A			Michelson	6,387,126	B1		Cumming
	4,718,904 A			Thornton	6,398,809			Hoffmann et al.
	4,738,680 A			Herman	6,406,494 6,413,262			Laguette et al. Saishin et al.
	4,781,719 <i>A</i> 4,787,903 <i>A</i>			Kelman Grendahl	6,413,277			Neuhann
	4,787,903 <i>1</i> 4,790,847 <i>1</i>			Woods	6,423,094			Sarfarazi
	4,842,601		/1989		6,425,917		7/2002	
	4,871,363 A			Kelman	6,428,574			Valunin et al.
	4,888,012			Horn et al.	6,432,246 6,443,984		8/2002 9/2002	Jahn et al.
	4,888,015 <i>A</i> 4,892,543 <i>A</i>			Domino Turley	6,443,985		9/2002	
	4,932,966 A			Christie et al.	6,451,056			Cumming
4	4,932,968 <i>A</i>	A 6		Caldwell et al.	6,461,384			Hoffmann et al.
	4,936,850 A			Barrett	6,464,725 6,475,240		10/2002 11/2002	
	4,944,082 <i>1</i> 4,950,288 <i>1</i>			Jones et al. Kelman	6,482,229			Gwon et al.
	4,963,148 A			Sulc et al.	6,488,707		12/2002	Callahan et al.
	5,104,590	A 4	/1992	Blake	6,488,708	B2		Sarfarazi
	5,108,429 A		/1992	Wiley	6,488,709		12/2002	
	5,185,107		/1993		6,494,911 6,503,276			Cumming Lang et al.
	RE34,424 I 5,282,851 <i>I</i>			Walman Jacob-LaBarre	6,517,577			Callahan et al.
	5,288,293 A			O'Donnell, Jr.	6,524,340	B2	2/2003	Israel
	5,306,297			Rheinish et al.	6,537,317			Steinert et al.
	5,326,347			Cumming	6,543,453 6,547,822			Klima et al.
	5,366,500 A			Schneider et al.	6,554,859		4/2003 4/2003	Lang et al.
	5,423,929 <i>I</i> 5,476,514 <i>I</i>			Doyle et al. Cumming	6,558,395			Hjertman et al.
	5,489,302			Skottun	6,558,420		5/2003	
:	5,496,366	A 3		Cumming	6,576,012		6/2003	
	5,507,806 A		/1996		6,592,621 6,596,025			Domino Portney
	5,527,415 <i>I</i> 5,549,668 <i>I</i>			Doyle et al. O'Donnell, Jr.	6,605,093		8/2003	
	5,562,731 A			Cumming	6,616,692		9/2003	Glick et al.
	5,571,177			Deacon et al.	6,638,306			Cumming
	5,578,081			McDonald	6,660,035			Lang et al. Callahan et al.
	5,607,472 A			Thompson	6,666,887 6,695,881			Peng et al.
	5,674,282 <i>I</i> 5,683,456 <i>I</i>		/1997 /1997	Cumming Blake	6,749,634		6/2004	Hanna
	5,702,441 A		/1997		6,755,859		6/2004	Hoffmann et al.
	5,728,155			Anello et al.	6,761,737		7/2004	Zadno-Azizi et al.
	5,728,156 A			Gupta et al.	6,764,511 6,767,363		7/2004 7/2004	Zadno-Azizi et al. Bandhauer et al.
	5,769,889 A 5,782,911 A			Kelman Herrick	6,786,928			Callahan et al.
	5,782,911 A			Lieberman	6,786,934		9/2004	Zadno-Azizi et al.
	5,803,925 A			Yang et al.	6,790,232		9/2004	Lang
:	5,843,187	A 12	/1998	Bayers	6,797,003 6,800,091			Blake et al. Callahan et al.
	5,855,605 A			Herrick	6,824,563		11/2004	
	5,964,802 <i>I</i> 5,984,962 <i>I</i>			Anello et al. Anello et al.	6,846,326	B2	1/2005	Zadno-Azizi et al.
	5,013,101 A		/2000		6,849,091			Cumming
	5,015,435 A			Valunin et al.	6,855,164			Glazier
	5,027,531			Tassignon	6,858,040 6,884,261			Nguyen et al. Zadno-Azizi et al.
	5,048,364 <i>E</i> 5,051,024 <i>E</i>			Skottun Cumming	6,899,732			Zadno-Azizi et al.
	5,051,024 <i>I</i>			Tran et al.	6,918,930		7/2005	Portney
	5,083,261			Callahan et al.	6,921,415			Callahan et al.
	5,096,077 <i>I</i>			Callahan et al.	6,921,416 6,926,736			Khoury
	5,096,078 <i>A</i>			McDonald Barraquer et al.	6,926,736			Peng et al. Bos et al.
	5,110,202 A 5,139,576 A			Doyle et al.	6,932,839			Kamerling et al.
	5,176,878 I			Gwon et al.	6,969,403	B2	11/2005	Peng et al.
(5,179,843 I	B1 1	/2001	Weiler	6,986,763			Holmen
	5,190,410 I			Lamielle et al.	6,986,787			Baker, Jr.
	5,197,058 I 5,197,059 I			Portney Cumming	6,991,651 7,025,783			Portney Brady et al.
	5,197,059 I 5,200,342 I			Tassignon	7,025,783		5/2006	Lindacher et al.
	5,200,342 I 5,200,344 I			Lamielle et al.	7,030,931		5/2006	Nagamoto
	5,217,612 I			Woods	7,041,134		5/2006	Nguyen et al.
(5,224,628 I	B1 5	/2001	Callahan et al.	7,048,759	B2	5/2006	Bogaert et al.

(56)	References Cit			ces Cited		8,034,108			Bumbalough
	U.S. PATENT			DOCUMENTS	8,038,711 8,043,370		10/2011 10/2011	Bretthauer et al.	
						8,043,372			Bumbalough
	7,048,760	B2		Cumming		8,048,155 8,048,156			Shadduck Geraghty et al.
	7,060,094			Shahinpoor et al.		8,052,752			Woods et al.
	7,063,723 7,087,080		6/2006 8/2006	Zadno-Azizi et al.		8,062,361			Nguyen et al.
	7,097,660			Portney		8,062,362			Brady et al.
	7,118,596			Zadno-Azizi et al.		8,066,768		11/2011	
	7,118,597		10/2006 10/2006	Miller et al.		8,066,769 8,070,806		11/2011 12/2011	
	7,122,053 7,125,422			Woods et al.		8,080,017		12/2011	
	7,137,702			Piers et al.		8,100,965			Cumming et al.
	7,150,759			Paul et al.		8,109,998 8,123,729			Cumming Yamamoto et al.
	7,150,760 7,192,444		12/2006	Zhang Blake et al.		8,123,729			Aharoni et al.
	7,192,444			Nguyen		8,158,712	B2	4/2012	Your
	7,204,849		4/2007	Portney		8,163,015		4/2012	Cumming
	7,217,288			Esch et al.		8,182,531 8,187,325			Hermans et al. Zadno-Azizi et al.
	7,220,279 7,226,478		5/2007 6/2007	Nun Ting et al.		8,206,442			Sel et al.
	7,229,475		6/2007			8,215,770	B2		Blum et al.
	7,238,201	B2		Portney et al.		8,216,305			Salvati et al. Coroneo
	7,247,168			Esch et al.		8,216,306 8,216,308			Blake et al.
	7,261,737 7,354,451		4/2008	Esch et al.		8,231,672			Deacon et al.
	7,377,640		5/2008	Piers et al.		8,235,525			Lesage et al.
	7,381,221			Lang et al.		8,241,353 8,241,355			Deacon et al. Brady et al.
	7,384,429 7,404,637		6/2008	Hanna Miller et al.		8,246,679			Nguyen et al.
	7,404,638			Miller et al.		8,267,996	B2	9/2012	Niwa et al.
	7,435,259		10/2008	Cumming		8,273,123			Ben Nun
	7,452,362			Zadno-Azizi et al.		8,303,656 8,314,927			Shadduck Choi et al.
	7,452,378 7,455,691			Zadno-Azizi et al. Feingold et al.		8,328,869			Smiley et al.
	7,462,193			Nagamoto		8,343,216		1/2013	Brady et al.
	7,485,144		2/2009			8,343,217			Bumbalough
	7,494,505 7,503,938			Kappelhof et al. Phillips		8,349,006 8,357,196			Zhao et al. Jain et al.
	7,553,327			Cumming		8,361,145	B2	1/2013	Scholl et al.
	7,569,073	B2	8/2009	Vaudant et al.		8,366,653			Shareef et al.
	7,591,849			Richardson		8,377,123 8,377,125		2/2013	Evans et al. Kellan
	7,601,169 7,615,056		10/2009	Ayton et al.		8,382,831			Ben Nun
	7,615,073			Deacon et al.		8,382,832			Deacon et al.
	7,621,949			Deacon et al.		8,398,709 8,403,984			Ben Nun Tsai et al.
	7,637,947 7,662,179			Smith et al. Sarfarazi		8,414,646			De Juan, Jr. et al.
	7,670,371			Piers et al.		8,425,595	B2	4/2013	Tsai et al.
	7,674,288	B2	3/2010	Nagamoto		8,425,597 8,425,598			Glick et al.
	7,713,299			Brady et al. Zadno-Azizi et al.		8,425,599			Klink et al. Shadduck
	7,744,646 7,744,647			Barrett		8,430,928		4/2013	Liao
	7,763,069	B2		Brady et al.		8,435,289			Cole et al.
	7,763,070			Cumming		8,449,611 8,454,688			Richardson Esch et al.
	7,771,471 7,780,729		8/2010 8/2010	Nguyen et al.		8,465,544			Brady et al.
	7,790,824			Freeman		8,475,527			Peterson et al.
	7,790,825			Lehman et al.		8,475,529 8,480,734		7/2013	Clarke Kellan et al.
	7,794,497 7,794,498			Brady et al. Pinchuk		8,486,140			Willis et al.
	7,794,498		10/2010			8,486,141		7/2013	Lang et al.
	7,815,678	B2		Ben Nun		8,486,142			Bumbalough et al.
	7,837,730			Cumming et al.		8,496,701 8,500,804			Hermans et al. Brady et al.
	7,842,087 7,854,764			Ben Nun Ben Nun		8,500,806		8/2013	
	7,857,850			Mentak et al.		8,523,942			Cumming
	7,871,437			Hermans et al.		8,529,623 8,535,376			Piers et al. Altmann
	7,878,655 7,883,540			Salvati et al. Niwa et al.		8,535,576			Woods et al.
	7,883,340			Altmann		8,551,164			Willis et al.
	7,931,686	B2	4/2011	Vaudant et al.		8,551,167	B2	10/2013	Cuevas
	7,981,155			Cumming		8,556,967			Sarfarazi
	7,985,253 7,998,199			Cumming Pan Nun		8,562,674 8,568,478			Cole et al. Zickler et al.
	8,012,204			Ben Nun Weinschenk, III et al.		8,574,293			Kappelhof et al.
	8,034,106			Mentak et al.		8,579,971		11/2013	
	8,034,107	B2	10/2011	Stenger		8,579,972	B2	11/2013	Rombach

(56)		Referen	ces Cited		2004/0111153			Woods et al.
ī	HS E	PATENT	DOCUMENTS		2004/0158322 2004/0162612		8/2004 8/2004	Portney et al.
·	0.0.1		DOCCINENTS		2004/0181279		9/2004	Nun
8,585,758		11/2013			2004/0215340 2004/0236422			Messner et al. Zhang et al.
8,585,759 8,608,799		11/2013	Bumbalough Blake		2004/0236423			Zhang et al.
8,608,800		12/2013			2004/0243233		12/2004	Phillips
8,613,766			Richardson et al.		2004/0249455 2005/0021137		1/2004	Tran Blake et al.
8,623,082 8,647,384		2/2014	Kappelhof et al.		2005/0021137		1/2005	
8,652,206			Masket		2005/0049700			Zadno-Azizi et al.
8,657,877			Glazier		2005/0055092 2005/0060032		3/2005	Nguyen et al. Magnante et al.
8,657,878 2001/0001836			Mentak et al. Cumming	623/6.37	2005/0071002			Glazier
2001/0012964			Lang et al.	025/0.57	2005/0085907		4/2005	
2001/0016771			Cumming		2005/0090896 2005/0096741			Ben Nun Cumming
2001/0044657 2001/0051826		11/2001	Rellan Bogaert et al.		2005/0000741		5/2005	
2002/0002404			Sarfarazi		2005/0107875			Cumming
2002/0016630		2/2002			2005/0113914 2005/0119739			Miller et al. Glazier
2002/0045937 2002/0072673			Sarfarazi Yamamoto et al.		2005/0119740			Esch et al.
2002/0101564			Herrick		2005/0125055			Deacon et al.
2002/0103535			Portney		2005/0125056 2005/0125057			Deacon et al. Cumming
2002/0107568 2002/0111678			Zadno-Azizi et al. Zadno-Azizi et al.		2005/0125059			Pinchuk et al.
2002/0111078			Ting et al.		2005/0131535		6/2005	
2002/0116058			Zadno-Azizi et al.		2005/0137703 2005/0143814		6/2005	Chen Esch et al.
2002/0116059 2002/0116060			Zadno-Azizi et al. Nguyen et al.		2005/0149184			Bogaert
2002/0116061			Zadno-Azizi et al.		2005/0209692	A1	9/2005	Zhang
2002/0116062			Portney		2005/0234547 2005/0251254			Nguyen et al. Brady et al.
2002/0133228 2002/0138140		9/2002 9/2002			2005/0251254			Nguyen et al.
2002/0138141			Zadno-Azizi et al.		2006/0020339	A1	1/2006	Ran
2002/0143395		10/2002			2006/0030938 2006/0041307			Altmann Esch et al.
2002/0161435 2002/0173847		10/2002	Portney Pham et al.		2006/0041307		3/2006	
2002/0177896		11/2002			2006/0064162	A1	3/2006	Klima
2002/0183843			Blake et al.		2006/0069433 2006/0089712		3/2006	Nun Malecaze
2002/0193876 2003/0018386			Lang et al. Laguette et al.		2006/0089712			Esch et al.
2003/0018380			Singer et al.		2006/0100703	A1	5/2006	Evans et al.
2003/0033013		2/2003	Callahan et al.		2006/0100704 2006/0116765			Blake et al. Blake et al.
2003/0060878 2003/0065387			Shadduck Callahan et al.		2006/0110703			Vaudant et al.
2003/0074060			Zadno-Azizi et al.		2006/0149369			Cumming et al.
2003/0074061			Pham et al.		2006/0178741 2006/0178742			Zadno-Azizi et al. Nagamoto
2003/0078655 2003/0078656			Callahan et al. Nguyen		2006/0178742			Nguyen et al.
2003/0078657		4/2003	Zadno-Azizi et al.		2006/0212116		9/2006	
2003/0078658			Zadno-Azizi		2006/0212117 2006/0247766		9/2006 11/2006	Lang et al.
2003/0083744 2003/0093149			Khoury Glazier		2006/0247767		11/2006	
2003/0105522			Glazier		2006/0253196		11/2006	
2003/0109926			Portney		2006/0259139 2006/0259140		11/2006	Zadno-Azizi et al.
2003/0114927 2003/0130732			Nagamoto Sarfarazi		2006/0271186			Nishi et al.
2003/0135271		7/2003	Bandhauer		2006/0271187			Zadno-Azizi et al.
2003/0135273			Callahan et al.		2007/0005135 2007/0005136			Makker et al. Richardson
2003/0149480 2003/0171808			Shadduck Phillips		2007/0010880	A1	1/2007	Esch
2003/0171809	A1	9/2003	Phillips		2007/0010882		1/2007	
2003/0187504 2003/0199976		10/2003 10/2003	Weinschenk, III et al.		2007/0016293 2007/0021831		1/2007 1/2007	Tran Clarke
2003/0199970			Peng et al.		2007/0021832	A1	1/2007	Nordan
2003/0204255	A1	10/2003	Peng et al.		2007/0027540			Zadno-Azizi et al.
2003/0204256 2004/0015236			Peng et al. Sarfarazi et al.		2007/0032866 2007/0050024		3/2007	Portney Zhang
2004/0013230			Heyman		2007/0050025			Nguyen et al.
2004/0064182	A1	4/2004	Kelman		2007/0067030		3/2007	Glazier et al.
2004/0073304 2004/0082993		4/2004 4/2004	Weinschenk, III et al.		2007/0078515 2007/0083260		4/2007 4/2007	•
2004/0082993			Woods et al.		2007/0083260			Colvard
2004/0082995		4/2004			2007/0088433			Esch et al.
2004/0106992			Lang et al.		2007/0100444		5/2007	Brady et al.
2004/0106993 2004/0111152			Portney Kelman		2007/0106377 2007/0106380		5/2007 5/2007	Smith et al. Terwee et al.
2004/0111132	AI	0/2004	Kellilali		2007/0100380	AI	3/2007	Terwee et al.

(56)		Referen	ices Cited	2010/0094415	A1		Bumbalough
		DATES	Doorn (D) TEO	2010/0121443			Michel et al.
	U.S.	PATENT	DOCUMENTS	2010/0121444 2010/0131059			Ben Nun Callahan et al.
2007/0106281	1 4 1	5/2007	D1-1	2010/0131039			Callahan et al.
2007/0106381 2007/0108643		5/2007 5/2007	Zadno-Azizi et al.	2010/0134754			Hong et al.
2007/0108045			Pynson	2010/0137983	A1		Culbertson et al.
2007/0123981		5/2007	Tassignon	2010/0152848			Williamson et al.
2007/0129798		6/2007	Chawdhary	2010/0179652			Yamamoto et al.
2007/0129803			Cumming et al.	2010/0198349 2010/0211167			Brady et al. Glazier
2007/0135915 2007/0142913		6/2007		2010/0211107		8/2010	
2007/0142913			Phillips Stenger	2010/0211171		8/2010	Sarfarazi
2007/0185574			Ben Nun	2010/0228260	A1		Callahan et al.
2007/0203578	3 A1		Scholl et al.	2010/0228344			Shadduck
2007/0213813	7 A1		Esch et al.	2010/0228346		9/2010	
2007/0239274		10/2007		2010/0292789 2010/0324672			Willis et al. Esch et al.
2007/0244560			Ossipov et al. Ben Nun	2010/0324673			Nguyen et al.
2007/0244561 2007/0260308		11/2007		2011/0029074			Reisin et al.
2007/0260309			Richardson	2011/0035001		2/2011	
2007/0260310			Richardson	2011/0035002		2/2011	
2008/0004699			Ben Nun	2011/0040376			Christie et al.
2008/0015689			Esch et al.	2011/0040378 2011/0040379			Werblin Bumbalough
2008/0021549 2008/0021550			Eagan et al. Richardson	2011/0040379		3/2011	Bumbalough
2008/0021330			Cole et al.	2011/0054601			Kadziauskas et al.
2008/0046074			Smith et al.	2011/0071628	A1		Gross et al.
2008/0046075			Esch et al.	2011/0082544			Ben Nun
2008/0051801			Hovey et al.	2011/0093067			Michalek et al.
2008/0077238			Deacon et al.	2011/0098810 2011/0098812			Altmann Ben Nun
2008/0077239 2008/0086208			Zickler et al. Nordan	2011/0098812			Ben Nun
2008/0086208		5/2008		2011/0112636			Ben Nun
2008/0109078			Rozakis et al.	2011/0112638	A1		Hermans et al.
2008/0154364			Richardson et al.	2011/0118836			Jain et al.
2008/0161913			Brady et al.	2011/0153014			Zhang et al.
2008/0161914			Brady et al.	2011/0153015 2011/0160852			Simonov et al. Mentak et al.
2008/0183289			Werblin	2011/0184514			Angelopoulos et al.
2008/0188930 2008/0208335			Mentak et al. Blum et al.	2011/0191086			Callahan et al.
2008/0208333			Werblin	2011/0238174	A1		Hong et al.
2008/0269882			Simpson et al.	2011/0245920			Richardson
2008/0269885			Simpson et al.	2011/0251686		10/2011	Masket Bumbalough et al.
2008/0269886			Simpson et al.	2011/0257742 2011/0270389			Glazer et al.
2008/0281417 2008/0300680		12/2008	Nagamoto	2011/0282441			Zadno-Azizi
2008/0306588			Smiley et al.	2011/0282442			Scholl et al.
2009/0005866			Cumming	2011/0282443			Smiley et al.
2009/0012609			Geraghty et al.	2011/0288638			Smiley et al.
2009/0018652			Hermans et al.	2011/0313522 2011/0313523		12/2011 12/2011	
2009/0033863 2009/0036982			Blum et al. Aharoni et al.	2012/0010704			Bumbalough
2009/0030982			Rombach	2012/0016473	$\mathbf{A}1$		Brady et al.
2009/0124773			Zhou et al.	2012/0029632	A1		Ben Nun
2009/0125105	5 A1		Lesage et al.	2012/0035724		2/2012	
2009/0125106			Weinschenk, III et al.	2012/0046743 2012/0046744			Pinchuk Woods et al.
2009/0171458 2009/0187242			Kellan et al. Weeber et al.	2012/0059465			Brady et al.
2009/0187242			Zhou et al.	2012/0078361			Shadduck
2009/0204209		8/2009		2012/0078363		3/2012	
2009/0204210			Pynson	2012/0078364			Stenger
2009/0204211			Angelopoulos et al.	2012/0116506 2012/0130487			Compertore Doraiswamy et al.
2009/0210054			Weeber et al.	2012/0130487			Doraiswamy et al.
2009/0228101 2009/0234448			Zadno-Azizi Weeber et al.	2012/0143327			Bumbalough
2009/0234449			De Juan, Jr. et al.	2012/0150292	A1	6/2012	Mentak et al.
2009/0248152			Bumbalough	2012/0203338		8/2012	
2009/0248154		10/2009		2012/0232648			Kahook et al.
2009/0264998			Mentak et al.	2012/0232650 2012/0232651			Hermans et al. Kahook et al.
2009/0264999 2009/0265000			Cumming Vaudant et al.	2012/0232631		9/2012	
2009/0283000			Gontijo et al.	2012/0243084			Geraghty et al.
2009/0292355			Boyd et al.	2012/0253459			Reich et al.
2009/0312836			Pinchuk et al.	2012/0290084		11/2012	
2009/0319040		12/2009	Khoury	2012/0296425			Cumming
2010/0016964			Werblin	2012/0296426			Brady et al.
2010/0036490			Deacon et al.	2012/0303119			Callahan et al.
2010/0094412	Z AI	4/2010	Wensrich	2012/0310338	ΑI	12/2012	Christie et al.

(56)	Referen	ices Cited	2013/0231741 A1	9/2013	Clarke
			2013/0231742 A1	9/2013	Deacon et al.
IJ.	S. PATENT	DOCUMENTS	2013/0238091 A1	9/2013	Danta et al.
			2013/0245754 A1	9/2013	Blum et al.
2012/0310342 A	1 12/2012	Nguyen et al.	2013/0245756 A1	9/2013	Liao
2012/0310342 A 2012/0310344 A			2013/0268070 A1	10/2013	Esch et al.
2012/0310345 A			2013/0268071 A1	10/2013	Vilupuru et al.
2012/0310343 A 2012/0323320 A		Simonov et al.	2013/0282117 A1	10/2013	Van Heugten et al.
2012/0323320 A 2012/0323321 A		Simonov et al.	2013/0297018 A1	11/2013	Brady et al.
2012/0323321 A 2012/0330415 A			2013/0304202 A1	11/2013	Basinger
2013/0013060 A		Zadno-Azizi et al.	2013/0304204 A1	11/2013	Bumbalough et al.
2013/0013000 A 2013/0018461 A			2013/0310931 A1	11/2013	
2013/0030525 A		Brady et al.	2013/0310932 A1	11/2013	Kellan
2013/0030323 A 2013/0035760 A			2013/0317606 A1	11/2013	Culbertson et al.
2013/0053760 A 2013/0053954 A			2013/0317608 A1	11/2013	Hermans et al.
2013/0060330 A		Weeber et al.	2013/0331937 A1	12/2013	Stevens
2013/0060330 A			2013/0338767 A1	12/2013	Mazzocchi et al.
2013/0073039 A			2013/0345713 A1	12/2013	Cole et al.
2013/0073035 A 2013/0103146 A			2014/0005780 A1	1/2014	Zhao
2013/0103140 A 2013/0103147 A		Christie et al.	2014/0005781 A1	1/2014	Zhao et al.
2013/0103147 A 2013/0116781 A		Ben Nun	2014/0005782 A1	1/2014	Kellan et al.
2013/0131794 A		Smiley et al.	2014/0052245 A1	2/2014	Zickler et al.
2013/0131754 A 2013/0138208 A			2014/0052246 A1	2/2014	Kahook et al.
2013/0150208 A 2013/0150961 A			2014/0058507 A1	2/2014	Reich et al.
2013/0166026 A		Bumbalough			
2013/0184816 A				CLICD DIE	DI ICATIONIC
2013/0190868 A		Kahook et al.	O.	THER PU	BLICATIONS
2013/0197635 A					
2013/0204364 A		Olson	Invitation to Pay Add	litional Fee	s mailed Sep. 17, 2013 in Interna-
2013/0204365 A		Dell	tional Application No	. PCT/US2	013/039708 (8 pages).
2013/0226294 A		Van Der Mooren et al.			· - ·
2013/0226295 A			* cited by examine	r	
2015/0220295 A	1 3/2013	De suan, st. et al.	chea by examine	•	




FIGURE 1





Figure 2






FIGURE 3




FIGURE 4

5c




FIGURE 5






FIGURE 6





FIGURE 7






Figure 9




FIGURE 10






Figure 1.







Figure 1






Figure 13

ACCOMMODATIVE INTRAOCULAR LENS AND METHOD OF IMPROVING ACCOMMODATION

CROSS REFERENCE TO RELATED APPLICATIONS

The present application is a U.S. Non-provisional application that claims the priority of U.S. provisional application No. 61/353,273, filed Jun. 10, 2010 and U.S. provisional application No. 61/368,862, filed Jul. 29, 2010. The contents of those applications are hereby incorporated by reference into the present disclosure in their entirety.

TECHNICAL FIELD

The present invention relates generally to intraocular lenses. More particularly, the present invention relates to accommodative, intraocular lens systems and methods for improving accommodation.

BACKGROUND INFORMATION

Under normal conditions, a healthy human eye focuses on near and distant objects by contraction and relaxation of the 25 ciliary muscle thereby contracting and releasing the tension on the zonules in the eye. The contraction of the ciliary muscle releases zonular tension (accomodative state) and allows the lens to alter to a more globular or spherical resting shape. The relaxation of the ciliary muscle increases tension 30 on zonules and elastic forces in the eye tissue overcome the inherent lens elasticity and result in stretching the lens equator and flattening the lens curvature (un-accomodative state).

In certain instances, for example when age-related opacification of the lens (cataract) interferes with vision, the natural crystalline lens of the eye needs to be removed. Generally, the natural lens is replaced with an artificial one, for example, an intraocular lens (IOL). Unfortunately, conventional IOLs, even those that profess to be accommodative, may be unable to provide sufficient spatial displacement of the lens along the 40 optical axis to provide an adequate amount of accommodation for near vision.

In conventional extracapsular cataract surgery, the crystalline lens matrix is removed by phacoemulsification through a curvilinear capsularhexis leaving intact the thin walls of the 45 anterior and posterior capsules, together with zonular ligament connections to the ciliary body and ciliary muscles. An intraocular lens is then placed in the capsular bag, which collapses around the IOL.

Conventional single-optic accommodative intraocular 50 lenses (AIOL) rely on the interaction of the ciliary muscle with the zonule and capsule to induce movement of the optic of the AIOL along its optical axis. Typically, the AIOL is secured within the capsular bag by two or more haptics that translate the radial stretching force exerted on the capsular 55 bag by the zonules in an attempt to achieve the desired axial displacement of the optic.

However, during the post-implantation fibrotic healing process, the anterior capsule fuses with the posterior capsule to form a rigid capsular disc. Loss of elasticity of the capsular 60 disc results and constrains the amount of movement that can be generated by the zonular force or elastic recoil of the intraocular lens and therefore, leads to a decrease in the amount of axial displacement of the lens that can be achieved.

Various lens systems have been designed to address this 65 loss of accommodation. Passive-shift single-optic lenses, the only accommodative lens currently marketed, were designed

2

to move forward under ciliary muscle contraction. Accommodation in these systems, however, remains limited by the loss of elasticity in the post-fibrotic capsule. Even the limited amount of accomodative amplitudes generated by these lenses immediately after surgery is lost within the first few weeks or month after surgery as capsular fibrosis ensures.

Accommodative lens designs with single or multiple optic lens assemblies have been disclosed, for example, in U.S. Pat. Nos. and U.S. application nos. 2009/0125106, 2005/0209692, 2007/0156236, 2009/0005866, 2007/0005136, 2009/0248154. Dual optic lenses retain the problem of capsular fibrosis and loss of amplitude/movement even though they are reported to provide a significant amount of accommodation. However, concerns about possible long-term formation of interlenticular opacification remain.

More recently, a lens systems that employs an active-shift mechanism using repulsive mini-magnets as a means of making accommodation partially independent of the zonules and mechanical properties of the capsular bag was disclosed (see U.S. Pat. Application Nos. 2009/0204210 and 2007/0118216. Still other methods of achieving accommodation include introduction of a polymerizable fluid with a desired refractive index into the capsular bag (lens refilling). Extensive investigation into the feasibility of these methods is still needed.

U.S. Publication No. 2009/0234449 discloses an intraocular lens comprising an accommodating element that is in contact with a substantial portion of the zonular region; the accommodating element is positioned relative to optical element and configured to cooperate with the ciliary muscle, the zonules and/or the vitreous pressure in the eye to effect a shape change to the optical element. According to the '449 publication, prior art multiple lens systems can be cumbersome and also require an axial displacement unachievable with a collapsed capsular bag and resulting ineffective accommodative mechanisms.

The need remains therefore, for an intraocular lens system and an effective mechanism for improving the accommodative capacity of an IOL following implantation. None of the current lens concepts take into account that the devitalized capsular bag after cataract surgery changes its physical properties from an elastic sphere to a contracted rigid disc.

SUMMARY OF THE INVENTION

Briefly, the present invention provides an intraocular lens system and method for improving accommodation that remedies the loss of axial and centrifugal movement caused by shrinkage and loss of flexibility of the capsular bag following implantation of conventional lens systems. A lens system of the present invention includes a flexible optic holder comprising a plurality of zonular capture haptics, in particular, regularly-spaced haptics that are adapted to allow or facilitate fusion of the capsular bag following placement of the optic holder within the capsular bag and ultimately to permit the sectioning of the fused capsular disc. During fusion, each haptic becomes permanently entrapped in its respective capsular disc section; sectioning frees adjacent haptics from each other; each haptic can, therefore, move independently in response to ciliary muscle and zonular forces on the capsule. The action of the zonular capture haptics can be translated to different types of optics suited to provide accomodative amplitude. The restored elasticity of the present accomodative IOL system, which allows the optic to return to a resting state when zonular tension is released, is provided by angulated haptics which straighten under zonular tension, by the elasticity of the optic or a combination of both.

The novel optic holder of the invention is implanted in two stages: first, the optic holder (which may or may not include an integral optic at this stage) is implanted and sufficient time is allowed for fusion of the anterior and posterior leaves of the capsular bag around the implanted device. In a second stage, the fused and fibrosed capsular bag is sectioned at regular intervals determined by spaces between the haptics of the optic holder, to reduce its rigidity, thereby restoring some of the movement lost during fusion of the capsule, fibrosis and formation of the capsular disc. The force of the zonules is uniformly transmitted to the lens via the entrapped haptics.

Unless it is already an integral part of the zonular capture haptic, the optic of the lens system may be inserted into the haptic during a second stage of the procedure. Any mechanical means or chemically-induced tensioning or positioning of the haptic/optic complex, which may have been employed to control the accomodative state of the haptic-lens complex during the fusion and contraction of the healing capsular bag would be removed at this juncture.

In one aspect, therefore, the present invention relates to a method for improving accommodation with an intraocular lens (IOL) system, the method comprising introducing a flexible optic holder with zonular capture haptics into the capsular bag of an eye of a subject and allowing a period of time 25 sufficient for fibrosis of the capsule to occur so that the zonular capture haptics are captured within the fused capsule. During this time the eye may be maintained in either an unaccommodative or accommodative state by administration of an agent to inhibit or induce accommodation, for example, 30 atropine or pilocarpine, respectively. Alternatively, mechanical means may be used to retain the capsule in a flattened and maximally (unaccommodated) or minimally (accommodated) expanded configuration. Apposition of the anterior and posterior leaves of the capsule to facilitate fusion may also be 35 achieved by introduction of an air bubble anterior to the capsular bag. Other mechanical, chemical or biological means may be utilized to enhance the adhesion of the anterior and posterior capsule.

In a second stage, cuts are made in the fibrotic capsular disc 40 at sectioning intervals between the zonular capture haptics of the optic holder. In some embodiments, during this second procedure, the optic is inserted into the holder.

In a related aspect, the invention relates to a flexible optic holder comprising an arrangement of zonular capture haptics 45 that permits 1) complete integration of the haptics during fusion of the capsule to form a capsular disc and 2) sectioning of the fused capsular disc. The optic holder comprises a plurality of hollow closed-loop or fenestrated haptics that extend outwardly from the center of the optic holder to define a disc that is roughly coextensive with the capsular disc and provides a template for sectioning of the disc following fusion. The optic holder of the invention is configured to provide support for an optic, to facilitate fusion and fibrosis of the capsular disc, to allow sectioning of the fused capsular 55 disc and to uniformly translate force from the zonules to the optic.

In one aspect, the invention relates to an intraocular lens system for implantation in a capsular bag of an eye, comprising 1) a flexible optic holder comprising a plurality of haptics 60 that extend outwardly from the center of the optic holder with sectioning regions therebetween where the haptics define a disc that is roughly coextensive with the capsular disc; and 2) an optic adapted to fit into the optic holder. The intraocular lens system is configured to be coextensive with the capsular 65 bag when placed therein and to become fixed within the capsular bag once fusion of the capsular bag has occurred.

4

The optic holder comprises a plurality of regularly spaced haptics, for example, from 3 to 120 haptics. In an alternate embodiment, the flexible optic holder comprises a single haptic with a fenestrated structure or of a surgical mesh or similar woven material with holes sufficiently large to allow contact and fusion of the anterior and posterior capsule through the fenestrae or holes. The haptics are spaced at regular intervals, with the space between adjacent members defining a sectioning region.

These and other objects, features and advantages of this invention will become apparent from the following detailed description of the various aspects of the invention taken in conjunction with the accompanying drawings.

BRIEF DESCRIPTION OF THE DRAWINGS

FIGS. 1a and 1b are illustrations of one embodiment of a flexible optic holder of the invention; FIG. 1b shows the optic holder in its expanded form.

FIGS. 2a and 2b show one embodiment of the invention inside the capsule of an eye prior to (2a) and following sectioning of the capsular bag (2b).

FIG. 3 is a schematic showing the position of the optic in the unaccommodated (top) and accommodated (bottom) states. During accommodation, the optic moves anteriorly (shown as downward); the dotted line indicates the position to which the optic returns when in the unaccommodated position.

FIG. 4a shows an alternate embodiment of the optic holder in the capsule, prior to sectioning, in which an optic has been secured. FIG. 4b shows the position of the haptics relative to the optic during accommodation; FIG. 4c shows the relative positions of the haptics and the optic under zonular tension (un-accommodative state). A staple-like retainer, for stabilizing the optic during fusion is also shown.

FIG. 5a shows an embodiment in which a snap-like fastener is used as optic attachment means for securing optic in optic holder; 5b shows the relative positions of the haptics and optic attachment means when the optic holder is fused within the capsule; 5c shows the relative positions of the haptics and optic during accommodation following sectioning of the capsular disc.

FIG. 6a shows anterior and posterior views of a dual-optic system in which two optic holders (black and gray) are used; FIG. 6b shows the position of the two optics during accommodation.

FIG. 7a shows an embodiment of the dual-optic system in which optics are an integral part of the optic holders; the positions of the two optics during accommodation (7b), and unaccommodation (7c) are shown.

FIG. 8 is an embodiment of an optic holder with a single dough-nut shaped haptic made of a woven or mesh-like material that can be cut, for example, with surgical scissors. A side view of the optic/optic holder is shown with a retainer, mechanical means for maintaining the minimum diameter (maximum contraction) of the capsular bag during fusion.

FIG. 9a shows an embodiment of the optic holder with an integrated flexible optic that approximates the natural lens; in the absence of zonular tension (accommodation), the optic is more spherical (9b) than when it is in the unaccommodated state (9c).

FIG. 10 is a photograph of an experimental model of a haptic design of the invention.

FIG. 11 includes a computer generated image and a schematic to illustrate the relative positions during accommoda-

tion of the haptics and a non-flexible optic using the optic holder of the present invention after the fused capsular bag has been sectioned.

FIG. 12 includes a computer generated image and a schematic to illustrate the relative positions of the haptics and 5 non-flexible optic in the unaccommodative state using the optic holder of the invention.

FIG. 13 includes a computer generated image to illustrate radial movement of the haptics when optic holder of the invention is used with a flexible (stretchable) optic.

DETAILED DESCRIPTION OF THE INVENTION

All patent applications, patents and other references cited herein are hereby incorporated by reference in their entirety 15 into the present disclosure.

In ophthalmology, the term "haptic" refers to a support structure that extends out from an optic element of an intraocular lens, for holding the lens in place within the capsular bag of the eye. For purposes of the present invention, 20 a haptic system designed to restore capsular flexibility lost "haptics" are sometimes referred to as "zonular capature haptics" and refer to structures or material that not only assist with placement and centration of the lens within the capsule, but are frame-like or fenestrated structures, which permit or facilitate fusion of the anterior and posterior capsules follow- 25 ing removal of the natural lens and placement of the artificial lens so that the haptics become securely entrapped within the fused capsule. The haptics define individual "sections" of the capsule which can be separated after fusion has occurred by making radial cuts in the capsule, specifically, beginning near 30 the center and extending out to the equator of the capsule. Following sectioning, each haptic is contained within a separate section of the capsule. The haptics of the optic holder of the invention have features which specifically enable it to become integrated into the fused capsule during fibrosis and 35 then uniformly translate the movement that results from contraction and relaxation of the ciliary body to the optic.

As used herein, the term "retainer" refers to a removable rigid device that is utilized to maintain the IOL system in controlled state of accommodation during the healing, fusion 40 period and is removed once fusion is complete. For example, IOL systems, which rely on angulated haptics as a mechanism of accommodation, would be maintained in an un-accommodated state during fusion of the capsular bag; keeping the optic holder in a flat planar configuration allows maximum 45 contact between the anterior and posterior capsule to enhance fusion and eliminate distortion of the capsular bag. In a single optic system, for example, the retainer may consist of a rigid rod with right angle extension inserting into each diametrically opposed pair of haptics. In a double optic system, a 50 simple suture ligation may be sufficient to maintain the two optics in apposition to each other, reducing the angle of the posterior and anterior haptics to a flat planar configuration. For flexible, prefilled or fillable pouches, the restraining device may be utilized to maintain the IOL system in an 55 accommodated position to encourage the contraction of the capsular bag to a minimum diameter, thereby maximizing the ability of the IOL system to stretch during relaxation of accommodation after the sectioning of the capsular bag.

Such a device may be mechanical or chemical, and may be 60 released by chemical, mechanical, laser or optical means.

As used herein, the term "fenestrated" indicates the presence of an opening or openings that allows for contact between the anterior and posterior capsule thereby facilitating fusion of the capsule through the opening(s).

Conventional accommodating lenses typically involve converting diametral movements of the ciliary muscle into

forward and backward movement of an optic portion of the IOL relative to the retina. For example, the only currently marketed accommodating IOL is a rigid, single optic IOL designed to rely on a forward translation of the optic to produce an increase in optical power of the eye. Movement of the IOL is produced by ciliary muscle contraction, capsular bag elasticity and/or suggested changes in vitreous cavity pressure to create an optical change in the eye.

However, implantation of the IOL into the capsule is fol-10 lowed by a natural physiological process not unlike applying a shrink-wrap film, in which the anterior and posterior capsular bag surfaces fuse around the haptics and seal the IOL within the fibrotic capsule. Furthermore, the fusing capsule undergoes fibrosis. During fibrosis the bag undergoes further contraction and loss of elasticity. As a result of this process, the IOL is immobilized within the fibrosed capsular disc and movement of the optic along the optical axis is extremely limited.

The present invention is directed to an optic holder having during fusion and fibrosis. This haptic system allows an implanted lens to transition more effectively between the accommodated and unaccommodated states, that is, in a fashion similar to the natural lens in response to forces applied to the capsule by the ciliary muscle and zonules. It achieves this by employing a haptic system that is flexible and becomes securely integrated into the fused capsular disc and allows for the capsular disc to be cut into sections, which has the effect of reducing the rigidity of the fused capsular disc and allows the optic holder to expand.

The present invention, therefore, is directed to a flexible optic holder comprising a plurality of haptics arranged in a circle and extending outwardly from a center portion of the optic holder (see FIGS. 1a and 1b). In addition to providing centration of the optic within the capsule like the haptics in conventional intraocular lenses, the haptic system of the optic holder of the present invention provide closed-loop, framelike structures that allow contact between the anterior and posterior capsules so that the process of capsular fusion and fibrosis are not impeded, thereby creating a skeletal support for the capsular disc. The natural post-phacoemulsification healing process is important for integration of the haptics into the capsular disc. Furthermore, the haptics are regularly arranged around the optic holder ring with a space between adjacent haptics to permit the fused capsular disc to be cut at regular intervals.

The haptics of the optic holder of the invention may be rigid or semi-rigid structures and may be made from a generally continuous element or a single continuous element of varying widths or thicknesses as long as the ability of the anterior and posterior capsules to securely fuse through the haptic is preserved. Haptics are made of a suitable nonabsorbable surgical material such as surgical wire, suture or the like. In one embodiment, haptics are constructed of polypropylene suture material, such as Prolene® (Ethicon, Somerville N.J.) The haptics of the optic holder may optionally include additional structures within the haptic frame, such as cross bars or anchors (for example, as shown in FIG. 1), to reinforce the haptic within the capsule following fusion. Anchors may be T-shaped, or a grid with cross-members that cross the length and/or width of the haptic.

The present invention also provides a two-stage process for inserting into an eye the intraocular lens system of the invention to achieve an improved level of accommodation. In one embodiment, evacuation of the capsular bag is followed by placement within the capsular bag of an optic holder that comprises one or more haptics that define a capsule-reinforc-

ing disc and which will ultimately receive the optic of the IOL system. Implantation of the haptic is followed by a healing interval, that is, a period of time sufficient to allow the anterior and posterior capsular bag surfaces to fuse together through and around the haptics of the optic holder thereby sealing the haptics within the fibrotic capsule. In one embodiment of the method, fusion of the capsular bag around the haptic occurs under conditions in which ciliary body movement is restricted, for example, by atropine-induced cycloplegia or pilocarpine-induced accommodation; paralysis of accommo- 10 dation movement optimizes capsular disc size, and enhances fibrosis of the capsule. Alternatively, mechanical means for maintaining the capsule in the unaccommodative or accommodative state, for example a retainer, may be employed to achieve the desired capsular disc size. The absence of an 15 accommodating optic during the phase of capsular fusion allows the optic holder to be free of mechanical strain and capsular bag distortion during the fusion and fibrosis period.

During the second stage the capsular disc and incorporated haptics are sectioned to reduce the rigidity of the capsular bag 20 so that the force exerted by the zonules can more effectively be transmitted to the capsule and permit movement of the optic along its optical axis and in the case of flexible optics, accommodative and unaccommodative movement caused by contraction of the ciliary muscle and/or zonular tension. Cuts 25 are made radially and at regular intervals between the haptics, (see FIG. 3) extending from the visual axis to the equator of the capsular disc. The accommodating optic may be inserted into the ring of the optic holder at the time of sectioning, either before or after sectioning. The optic contains means for securing the optic into the lens holder, for example a circumferential releasable connecting rib or series of releasable connecting tabs, pins, plugs or the like that snap into a corresponding receptacle: a groove, notch or hole on the inner edge of the haptic.

The intraocular lens system of the invention comprises a rigid or flexible optic, single lens or multiple lenses, or fillable or pre-filled, and in one embodiment, an accommodating optic and optic holder are inserted as a single unit at the time of initial cataract surgery. Such an integrated intraocular lens 40 system may include a restricting device to mechanically or chemically maintain the system in a specific state of accommodation during the fusion and fibrosis of the capsular disc, for example a maximally accommodated optic and pilocarpine-induced pharmacologic accommodation during the 45 entire duration of capsular fibrosis and fusion. This eliminates mechanical strain or movement or distortion of the capsular bag during the fusion/fibrosis phase after the first stage procedure, optimizing the size of the fibrotic capsular disc, sealing of the haptic members in the capsular disc and resting 50 tension on the zonlues.

During the second stage, radial cuts are made in the capsular disc, between the haptics, and extending from the visual axis to the outer edge of the capsular disc. This releases the restraining effect of fibrosis on the now segmented, capsular 55 bag. An optic restraining device, if employed, is also removed at this time allowing the accommodating optic to respond to the zonular tension transmitted by the haptics during relaxation of accommodation, or to return to its resting accommodating state during contraction of ciliary body and relaxation of the zonules.

The method of the present invention for the implantation of an intraocular lens, therefore, includes making a plurality of regularly-spaced radial cuts around the capsule/haptics, extending from the visual axis to the edge of the capsular bag. 65 Sectioning alters the rigidity of the capsule following capsular fibrosis/fusion so that the tension and relaxation of the

8

zonules is more effectively translated to the capsule thereby enhancing spatial displacement of the optical element.

Uniform and complete natural, therapeutically-assisted or -enhanced fusion of the capsule around the haptic prior to optic placement ensures uniformity of sectioning essential to centration and stability of the optic once it is inserted.

Following insertion, the patient undergoes a recovery period of one to six weeks, preferably two to three weeks, for complete fibrosis of the capsule to occur. A second surgical procedure is performed to restore "elasticity" to the capsule by making radial incisions in the capsule at regular intervals forming roughly triangular or trapezoidal sections, each of which contains one of the haptics of the optic holder.

By "sectioning" the capsular disc, greater movement of the optic along the optical axis is achieved. Each haptic-reinforced section of the sectioned disc is generally separate from the others, held together by the inter-haptic connectors of the lens holder and the optic, when it is positioned within the optic holder. As a result of the sectioning, however, the force generated by the zonules is more effectively transmitted to the optic.

It is hypothesized that a 1 mm change in capsular bag diameter results in 2 mm anterior/posterior optic movement. A 12 mm capsular bag under zonular tension, (i.e. unaccommodated) collapsing to 11 mm in diameter as a result of the AIOL tension will result in a 2 mm anterior displacement during accommodation. Expected ranges of total accommodation with rigid single and double optics and flexible deformable optics are shown in Table 1.

TABLE 1

	Single Optic	Dual Optic	Flexible Optic
Actual accommodation	2D	6D	4-7D
Pseudo accommodation	1.5-2D	1.5-2D	1.5-2D
Total accommodation	3.5-4D	7.5-8D	4.5-9D

Intraocular Lens System

The intraocular lens system of the invention comprises 1) a flexible optic holder specifically adapted to permit fusion of the capsule and sectioning of the capsular disc once fusion is complete and 2) an optic. The optic holder comprises a plurality of haptics adapted to receive and secure the optic(s). The optic holder supports the optic around its equator and couples the optic to the capsular bag of the eye. Following extraction of the natural lens and placement of the flexible optic holder of the invention into the capsular bag, the natural healing process causes the optic holder to become entrapped, "capturing" the capsule, when the anterior and posterior leaves of the capsular bag fuse together. Subsequently, radial cuts to the capsule allow the sections of the fused capsule and the captured haptics to move independently of the others in response to zonular tension.

Optic Holder with Zonular Capture Haptics

The intraocular lens of the invention comprises a optic holder with zonular capture haptics, that is, haptics, which by their closed-loop design, allow maximal fusion of the anterior and posterior capsule through the haptics and permit the capsule to be sectioned following fusion, so that, in response to tension by the zonules, each haptic is able to move radially away from the center of the optic holder and then return to the initial position when zonular tension is released. The haptics

become fused within the capsular bag and enables the capsule to be sectioned into a plurality of capsular "sections." The haptics provide fixation, centration and stability of the optic (s) within the eye and provide a skeletal support for the capsular bag so that its rigidity can be reduced by cutting it of the capsular bag so that its rigidity can be reduced by cutting it of the capsular bag so that its rigidity can be reduced by cutting it of the capsular bag so that its rigidity can be reduced by cutting it of the capsular bag so that its rigidity can be reduced by cutting it of the capsular bag so that its rigidity can be reduced by cutting it of the capsular bag so that its rigidity can be reduced by cutting it of the capsular bag so that its rigidity can be reduced by cutting it of the capsular bag so that its rigidity can be reduced by cutting it of the capsular bag so that its rigidity can be reduced by cutting it.

One embodiment of an optic holder of the present invention is shown in FIG. 1. Optic holder 10 comprises a plurality of haptics 20 extending outwardly from the center of optic holder 10. The arrangement of haptics 20 generally defines a ring that receives an optic 80, and haptics 20 further comprise optic attachment means 50/52 to hold the optic 80 securely in place. In this embodiment, haptics 20 are roughly trapezoidal in shape. Adjacent haptics 20 are connected via an interhaptic loop or connector 30 of flexible material so that the inner edges of haptics 20 form a continuous ring to which the optic is ultimately attached.

In some embodiments, optic holder 10 is flexible (FIG. 1b). The haptics 20 are spaced apart at regular intervals to generally form a disc that is roughly coextensive in size with the capsular bag and retains the circular shape of the fused capsular disc. Following placement of an optic holder of the invention into the capsular bag, the capsule will shrink and fuse around the optic holder (much like shrink-wrap). As 25 shown in FIG. 2b, once fusion of the capsule is complete, cuts are made between the haptics of the optic holder. This allows the sections to move somewhat independently of each other.

FIG. 2a shows an embodiment of an optic holder of the invention within the unsectioned capsular bag (shaded area). 30 Following sectioning between haptics (FIG. 2b), the optic experiences improved freedom of movement, compared to the unsectioned capsule in response to zonular tension.

In FIG. 3, the positions of the haptics and optic of any intraocular lens, including conventional intraocular lenses, 35 once the capsule has fused and become fibrosed are shown in the upper panel. The bottom panel shows anterior movement of the optic (shown as downward movement) made possible by sectioning the fused capsule using the optic holder of the present invention.

FIG. 4a shows an embodiment of an optic holder 100 in which an optic 180 has been secured prior to sectioning of the capsular bag; sectioning lines 184 are indicated. The position of the optic 180 and haptics 120 during accommodation (4b) and the unaccommodative state (4c) are shown.

FIG. 5 shows an embodiment of optic attachment means 52/50, for example, a pin 52 and receptacle 50 snap-like fastener, by which optic 80 is secured in optic holder. Because the haptics are angulated, without any other force in play, insertion of the optic into the holder forces the haptic into a 50 roughly 30° angle with the optic.

In one embodiment (see FIGS. 6 and 7), a pair of anteriorly- and posteriorly-angled optic holders are used in an alternating arrangement or configuration. The anterior haptic will receive an anterior optic, while the posterior one will 55 receive a posterior optic. In the dual-optic system, during accommodation, zonular tension is released and the haptic system returns to its resting state of maximum angulation of the optic attachment means causing the two lenses to move axially away from each other, thereby providing increased 60 accommodative amplitude.

When tension is applied to the optic holder during relaxation of ciliary body, the haptics straighten thereby causing an axial displacement of the optics towards each other (as shown in FIG. 7c). With a dual-optic system, maintenance of the 65 appositional relationship of anterior and posterior optics is important to ensure uniform fusion of the capsule. This may

10

be achieved mechanically or through the use of therapeutic agents that induce unaccommodation.

Referring to FIG. 8, in one embodiment, the optic holder 300 comprises an optic 380 or optic holding member at its center and a single doughnut-shaped haptic 320 extends outwardly from the optic 380 and is made of a woven material or mesh or fenestrated material that 1) permits fusion of the capsular bag and 2) can be sectioned or cut. Radial sectioning lines (not shown) may be indicated on the capsule-capture haptic and small holes near the optic/optic holding member provided for starting the cut.

FIG. 9 shows an embodiment in which haptics 420 and optic 480 comprise a single unit, with a flexible optic 480, or fillable or prefilled optic pouch attached directly to the haptics 420. Such an optic has a predetermined dioptric power in its resting state (9b). The inherent optic elasticity allows stretching of the optic equator and flattening of the lens curvature in the un-accomodative state (9c) and return to a more spherical shape (9b) during accommodation when zonular tension is released.

Flexible optics, fillable or prefilled optic pouchs are known in the art and can be integrated or adapted for use with the optic holder of the invention. The adaptation of any elastic optic embodiments to a zonular capture optic holder would enable the optic to change shape and accommodate.

In some embodiments, the haptics further comprise an anchor or other support structure for promoting fusion/fibrosis of the capsular bag and integration of the haptic within the capsular bag. The anchor may be any shape, for example, T-shaped, or size which will have the effect of securing and reinforcing the haptic within the capsular bag. Implantation of the Optic Holder

In an initial procedure, the optic holder is placed in the capsular bag of the eye, and the anterior and posterior leaves of the capsular bag are allowed to fuse together securing the optic holder within the fused capsular bag or disc. Uniform healing of capsule around optic holder ensures centration of the optic once it is placed in optic holder.

In a second procedure, a number of cuts are made in the capsular disc between the haptics of the optic holder. The cuts extend from the visual axis to the outer edge of the capsular disc. Additionally, the posterior zonules may become stiff further limiting anterior/posterior movement. In some situations, it may be desirable to cut the posterior zonules.

In one embodiment, the optic holder comprises a ring structure in which at least three capsule-capture members (haptics) extend outwardly from the center of the ring (see FIG. 1). The number and size of capsule-capture members of the optic holder varies depending on the number of sections which the clinician determines to be optimal. In so determining, an optimal number of sections may be determined to be that number which will permit the greatest axial movement of the optic that can be achieved without compromising the integrity of the capsular bag. Further considerations regarding the number of sections to be made include allocating an amount of time for sectioning which the clinician feels is appropriate for the safety and well-being of the patient.

The haptics of the optic holder are made from inert or biocompatible materials known to those of skill in the art, for example, silicone, polypropylene, acrylic polymers or the like. Haptics are made in an open configuration (loops), and may be of any shape, for example, generally triangular or trapezoidal, which, as a group roughly define a disc that is coextensive with the capsular bag. Generally, the outer edge of the haptics extend to the equator of the capsular bag to form a capsule-reinforcing disc which is roughly coextensive with the capsular disc, and by virtue of their shape or porosity,

permit fusion of the anterior and posterior leaves of the capsular bag to form a capsular disc encasing the haptic.

The number and configuration of individual members are chosen in accordance with considerations discussed supra to anchor the optic-holder ring in the capsular bag and form an internal "frame" on which the fused capsular disc is supported. To maximize or minimize the diameter of capsular bag during fusion thereby impacting the ultimate size of the fused capsular disc, accommodation can be controlled during fusion of the capsular bag by administration of an agent to inhibit or induce accommodation.

Optics

The intraocular lens system of the invention may comprise multiple lens holders and therefore, be able to hold one or more optics. Optics are generally symmetrical about the optical axis. Examples of suitable optics are well known in the art and can be adapted for use with the optic holder and zonular capture haptics of the invention. These include optics that are flexible, deformable, foldable, or rigid, preformed or fillable 20 and which are made from a liquid, solid or semi-solid material. In one embodiment, if a flexible optic is used, it can assist in accommodation not only by anterior-posterior displacement, but also by changing its radius of curvature.

Examples of suitable optic materials include silicone(s), 25 acrylics, hydrogels and other inert or biocompatible polymers known to those of skill in the art. In one embodiment, the optic comprises a means for securing the optic into the haptic, for example a circumferential releasable connecting rib or series of releasable connecting tabs or pins that snap into a corresponding groove, notch or hole on the haptic.

Haptics may incorporate any means suitable for attaching and securing the optic; these are well known in the art.

Sectioning of capsular disc after fusion and fibrosis of the haptic therein can be accomplished by virtue of a small gauge (for example, 23 or 25 gauge) trans-conjunctival vitrectomy system with trocars and cannulas placed diametrically opposed to the section line in the capsule/haptic where the section is to be made. A small gauge (for example, 23 or 25 gauge) scissor is introduced through a cannula and used to cut the capsule from the visual axis to the outer edge of the capsular disc.

While several aspects of the present invention have been described and depicted herein, alternative aspects may be effected by those skilled in the art to accomplish the same 45 objectives. Accordingly, it is intended by the appended claims to cover all such alternative aspects as fall within the true spirit and scope of the invention.

A haptic arrangement was manufactured from 4-0 surgical Prolene sutures with 7-0 surgical Prolene retention bands 50 (FIG. 10). The function of the retention bands are to maintain the shape of each haptic loop, to function as an additional retaining anchor within each haptic and to maintain the distance between the haptics during the sterilization process. The surgical procedure for implantation of an optic holder of 55 the invention consists of two stages.

All procedures related to the use of animals conform to the Guide for the Care and Use of Laboratory Animals (National Institutes of Health, National Research Council) and are approved by the Wisconsin National Primate Research Center. The animals are housed in facilities accredited by the Association for Assessment and Accreditation of Laboratory Animal Care International (AAALAC). Animal subjects are paired or individually housed. Diet consists of monkey chow supplemented with fresh fruit and vegetables. Water is freely 65 available. All animals are observed daily by a veterinary technical staff and caretakers for signs of ill health.

12

Adult Rhesus monkeys (Macacca mulatta) of either sex without any ocular disease, are the subjects of the following experiments.

At Stage 1, the treated eye is rendered aniridic by complete surgical removal of the iris. The absence of the iris facilitates evaluation of the optic holder's performance. Standard cataract removal by small, clear corneal incision and phacoemulsification is followed by implantation of the lens holder of the invention via standard 2 mm lens injector (Photo 2). The treated eye is maintained in a pharmacologic state of forced accommodation by the administration of pilocarpine drops 4 times per day. This allows the zonules to be free of tension during the healing, fibrosis, contracture phase of the capsular bag. A minimum capsular disc size is thus obtained, which is anticipated to securely incorporate the haptics of the lens holder in between the fused anterior and posterior capsules. Once the fibrosis phase is complete, approximately 2-4 weeks after the Stage 1 surgery, the animal is anesthetized again for the Stage 2 surgery.

A 23 or 25 gauge standard vitrectomy instrument is employed. Four trans-conjunctival cannulas are inserted at the standard pars plana location, coinciding with the planned capsular section lines as defined by the location of the spaces between the haptics. A central posterior capsulotomy is performed with the vitrectomy instrument. A 23 or 25 gauge Vitreoretinal scissor is introduced via each cannula to perform radial cuts extending from the edge of the posterior capsulotomy to the equator of the capsular bag, cutting across the fused capsular sheets, the 7-0 Prolene suture extending in between capsular members, to the edge of the capsular disc, ensuring that no connection remains between the individual sections of capsular disc capsule. The only connection between the sectioned capsules and entrapped haptics is the flexible inter-haptic loop extending from one haptic to the next.

Administration of pilocarpine drops is discontinued postoperatively. Once the eye has recovered from the surgical intervention, the eye is challenged with pharmacologic accommodation and relaxation of accommodation under anesthesia with short acting pharmacologic agents while the eye is monitored and videographed. It is anticipated that during relaxation of accommodation, zonular tension is produced and transmitted to the individual segments of the former capsular disc, which now move independently. Each haptic of the optic holder should move centrifugally and away from each other. When accommodation is induced pharmacologically, the tension of the zonules is released and the elasticity of the inter-haptic loops returns haptics to a closer configuration. A change in diameter of the optic holder of up to 1 mm is expected, based on previously published data.

I claim:

- 1. A method for improving accommodation of an intraocular lens system in an eye, the method comprising:
 - (a) providing the intraocular lens system comprising a plurality of spaced radial haptics;
 - (b) positioning at least a portion of the system in a capsular bag of the eye;
 - (c) allowing anterior and posterior leaves of the capsular bag to fuse; and
 - (d) making a plurality of substantially radial cuts in the fused anterior and posterior leaves of the capsular bag to produce sections each containing a haptic.
- The method of claim 1, wherein the system comprises an
 optic and an optic holder, the method further comprising inserting the optic into the optic holder after the capsular bag has fused.

- 3. The method of claim 1, wherein said intraocular lens system comprises an optic holder comprising the plurality of radial haptics.
- **4**. The method of claim **1**, wherein said plurality of haptics are regularly spaced and define sectioning intervals therebetween for the substantially radial cuts.
- 5. The method of claim 1, further comprising maintaining the eye in an unaccommodative state during the fusion of the capsular bag.
- **6**. The method of claim **1**, further comprising maintaining the eye in an accommodative state during the fusion of the capsular bag.
- 7. The method of claim 5, wherein maintaining the unaccommodative state comprises administering an agent to inhibit accommodation.
- 8. The method of claim 7, wherein said agent comprises atropine.
- 9. The method of claim 6, wherein maintaining the accommodative state comprises administering an agent to promote accommodation.
- 10. The method of claim 9, wherein said agent comprises 20 pilocarpine.
- 11. The method of claim 1, wherein the capsular bag fuses within about 1 to about 6 weeks.
- 12. The method of claim 1, wherein the capsular bag fuses within about 2 to about 4 weeks.

14

- 13. The method of claim 1, wherein said cuts extend from a visual axis to an edge of the fused capsular bag.
- 14. The method of claim 1, wherein the system comprises an optic and an optic holder, and wherein both the optic and the optic holder are positioned in the capsular bag prior to fusion.
- 15. The method of claim 1, wherein the system comprises an optic and an optic holder, the method further comprising releasably securing the optic in the optic holder.
- 16. The method of claim 1, wherein said plurality of haptics comprises no more than 120 haptics.
- 17. The method of claim 1, wherein the system further comprises a restricting device configured to maintain the haptics in a specific state of accommodation during fusion of the capsular bag.
- 18. The method of claim 1, wherein the system further comprises a retainer adapted to achieve a desired capsular disc size
- 19. The method of claim 1, wherein the plurality of substantially radial cuts comprises a plurality of single cuts between the plurality of radial haptics.
 - 20. The method of claim 1, wherein each haptic comprises a closed-loop structure.

* * * * *